Technical Data Sheet

Purified Rat Anti-Mouse CD4

Product Information

Material Number:
Alternate Name:
Size:
Concentration:
Clone:
Immunogen:
Isotype:
Reactivity:
Storage Buffer:

550280

Cd4; CD4 antigen; L3T4; Ly-4; T-cell surface antigen T4/Leu-3 1.0 ml 62.5 µg/ml RM4-5 Mouse Thymocytes (BALB/c) Rat (DA) IgG2a, κ QC Testing: Mouse Aqueous buffered solution containing BSA, goat serum, and ≤0.09% sodium azide.

Description

The RM4-5 monoclonal antibody specifically binds to the CD4 (L3T4) differentiation antigen expressed on most thymocytes, subpopulations of mature T lymphocytes (i.e., MHC class II-restricted T cells, including most T helper cells and immunosuppressive regulatory T cells), and a subset of NK-T cells. CD4 has also been reported to be detected on pluripotent hematopoietic stem cells, bone marrow myeloid and B-lymphocyte precursors, intrathymic lymphoid precursors, and a subset of splenic dendritic cells. CD4 has been reported to be expressed on the plasma membrane of mouse egg cells and is involved in adhesion of the egg to MHC class II-bearing sperm. CD4 is an antigen coreceptor on the T-cell surface which interacts with MHC class II molecules on antigen-presenting cells. It participates in T-cell activation through its association with the T-cell receptor complex and protein tyrosine kinase lck. Purified RM4-5 mAb has been reported to block the binding of FITC-conjugated anti-mouse CD4 clones GK1.5 and H129.19, but not the RM4-4 clone.



Immunohistochemical staining of mouse CD4. Frozen sections of normal mouse spleen were stained with the anti-mouse CD4 antibody clone RM4-5. CD4+ T lymphocytes can be identified by the intense brown labeling of their cell surface membranes. Amplification 20X.

Preparation and Storage

Store undiluted at 4°C.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Application Notes

Application				
Flow cytometry	Routinely Tested			
Immunohistochemistry-frozen	Tested During Development			
Immunohistochemistry-zinc-fixed	Tested During Development			
Immunohistochemistry-formalin (antigen retrieval required)	Not Recommended			

Recommended Assay Procedure:

Immunohistochemistry: Clone RM4-5 is recommended to test for immunohistochemical staining of acetone-fixed frozen or zinc-fixed paraffin embedded sections of mouse spleen or thymus. **IHC of formalin-fixed paraffin embedded sections is not recommended.** The isotype control recommended for use with this antibody is purified rat IgG2a (Cat. No. 559073). For optimal indirect immunohistochemical staining, the RM4-5 antibody should be titrated (1:10 to 1:50 dilution) and visualized via a three step staining procedure in combination with biotinylated polyclonal anti-rat Ig (Cat.No. 559286) as the secondary antibody and streptavidin-HRP (Cat.No. 550946) together with the DAB detection system (Cat.No.

BD Biosciences

bdbiosciences.	com				
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
6//.252.8995	800.979.9408	32.53.720.550	0120.8555.90	05.0001.0633	55.11.5165.9995
For country co	ntact informatio	on, visit bdbiosci	ences.com/conta	ct	
Conditions: The ir of any patents. Bl use of our produc product or as a co written authoriza	nformation disclose D Biosciences will n ts. Purchase does n Imponent of anoth tion of Becton. Dic	d herein is not to b ot be help responsi ot include or carry er product. Any use kinson and Compar	e constructed as a ro ble for patent infrin any right to resell ou of this product oth by is stictly prohibite	ecommendation to u gement or other vio r transfer this produc er than the permitte d.	use the above product in violatio lations that may occur with the ct either as a stand-alone ed use without the express
For Research Use	Only. Not for use in	diagnostic or there	apeutic procedures.	Not for resale.	
Unless otherwise	noted, BD, BD Logo	and all other trad	emarks are property	of Becton, Dickinso	n and Company. © 2011 BD



550880). More conveniently, the anti-rat Ig HRP detection kit (Cat. No. 551013) can be used which contains the biotinylated secondary antibody, the antibody diluent, Streptavidin-HRP and DAB substrates for use in the staining procedure.

Suggested Companion Products

Catalog Number Name		Size	Clone	
559073	Purified Rat IgG2a κ Isotype Control	0.25 mg	R35-95	
559286	Biotin Goat Anti-Rat Ig	0.5 mg	Polyclonal	
551013	Anti-Rat Ig HRP Detection Kit	200 tests	(none)	
550946	Streptavidin HRP	50 ml	(none)	
550880	DAB Substrate Kit	500 tests	(none)	
559148	Antibody Diluent for IHC	125 ml	(none)	

Product Notices

Since applications vary, each investigator should titrate the reagent to obtain optimal results. 1.

- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before 2. discarding to avoid accumulation of potentially explosive deposits in plumbing.
- Source of all serum proteins is from USDA inspected abattoirs located in the United States. 3.
- This antibody has been developed for the immunohistochemistry application. However, a routine immunohistochemistry test is not 4. performed on every lot. Researchers are encouraged to titrate the reagent for optimal performance.
- 5. An isotype control should be used at the same concentration as the antibody of interest.
- 6. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Bierer BE, Sleckman BP, Ratnofsky SE, Burakoff SJ. The biologic roles of CD2, CD4, and CD8 in T-cell activation. Annu Rev Immunol. 1989; 7:579-599. (Biology) Bosselut R, Zhang W, Ashe JM, Kopacz JL, Samelson LE, Singer A. Association of the adaptor molecule LAT with CD4 and CD8 coreceptors identifies a new coreceptor function in T cell receptor signal transduction. J Exp Med. 1999; 190(10):1517-1526. (Biology: Immunoprecipitation)

Frederickson GG, Basch RS. L3T4 antigen expression by hemopoietic precursor cells. J Exp Med. 1989; 169(4):1473-1478. (Biology)

Janeway CA Jr. The T cell receptor as a multicomponent signalling machine: CD4/CD8 coreceptors and CD45 in T cell activation. Annu Rev Immunol. 1992; 10:645-674. (Biology)

Nakamura T. Personal Communication. . (Immunogen: Blocking)

Wineman JP, Gilmore GL, Gritzmacher C, Torbett BE, Muller-Sieburg CE. CD4 is expressed on murine pluripotent hematopoietic stem cells. Blood. 1992; 180(7):1717-1724. (Biology)

Wu L, Antica M, Johnson GR, Scollay R, Shortman K. Developmental potential of the earliest precursor cells from the adult mouse thymus. J Exp Med. 1991; 174(6):1617-1627. (Biology)

Wu L, Scollay R, Egerton M, Pearse M, Spangrude GJ, Shortman K. CD4 expressed on earliest T-lineage precursor cells in the adult murine thymus. Nature. 1991; 349(6304):71-74. (Biology)

BD Biosciences

bdbiosciences.com United States Canada 877.232.8995

Europe Japan 800.979.9408 32.53.720.550 0120.8555.90 For country contact information, visit bdbiosciences.com/contact

Latin America/Caribbean 65.6861.0633 55.11.5185.9995



Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Buckness will not be help responsible for patent infringement or other violations that may occur with the product or as a component of another product. Any use of this product other than the permitted use without the express

Asia Pacific

written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD