

Technical Data Sheet

Purified NA/LE Mouse Anti-Human CD178


Product Information

Material Number:	556375
Alternate Name:	Fas Ligand, CD95 Ligand
Size:	0.25 mg
Concentration:	1.0 mg/ml
Clone:	NOK-2
Immunogen:	L5178Y Mouse T Lymphoma Cells Expressing Recombinant Human FasL
Isotype:	Mouse IgG2a
Reactivity:	QC Testing: Human
Target MW:	40, 26 kDa
Storage Buffer:	No azide/low endotoxin: Aqueous buffered solution containing no preservative, 0.2µm sterile filtered. Endotoxin level is ≤0.01 EU/µg (≤0.001 ng/µg) of protein as determined by the LAL assay.

Description

Fas (APO-1, CD95) is a 45 kDa cell surface protein that mediates apoptosis when cross-linked with agonistic anti-Fas antibodies or Fas ligand (FasL). Fas belongs to the TNF (tumor necrosis factor)/NGF (nerve growth factor) receptor family, and is expressed in various tissue and cells including the thymus, liver, ovary, and lung. FasL is a 40kDa TNF family membrane protein that induces apoptosis by binding to Fas, its cell-surface receptor. FasL is expressed on activated T and NK cells. Both Fas and FasL are thought to play an important role in the apoptotic processes that take place during T cell development.

NOK-2 recognizes human FasL. It recognizes both the membrane bound (FasL) and soluble (sFasL) forms. L5178Y mouse T lymphoma cells expressing recombinant human FasL were used as immunogen. FasL and sFasL migrate at reduced molecular weights of 40 and 26 kDa, respectively. However, the molecular weights observed in a particular sample may vary according to FasL and sFasL glycosylation and breakdown patterns as described in Tanaka et al.



Immunoprecipitation of Fas Ligand (FasL). T lymphoma L5187Y cells were transfected with human FasL cDNA and either treated with metalloprotease inhibitor KB8301 or left untreated. KB8301 blocks FasL cleavage resulting in high levels of cell surface FasL. FasL is detected in [35S]-labeled cells treated with inhibitor KB8301 (lane 1), but not in untreated cells (lane 3). Isotype (negative) controls are shown in lanes 2 and 4. Purified NOK-2 (Cat. No. 556376) or Mouse IgG2a (Cat. No. 553454) was used for immunoprecipitations.

Preparation and Storage

Store undiluted at 4°C.

This preparation contains no preservatives, thus it should be handled under aseptic conditions.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Application Notes

Application

Immunoprecipitation	Routinely Tested
Flow cytometry	Tested During Development
ELISA	Reported
Neutralization	Reported
Western blot	Not Recommended

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



Recommended Assay Procedure:

IP: Applications include immunoprecipitation (1-2 µg/one million cells).

Functional Assay: NOK-2 has also been shown to neutralize the cytotoxic activity of FasL. Neutralization of FasL activity inhibits Fas-mediated killing. The NA/LE format of NOK-2 (Cat. No. 556375) should be used for all functional assays. NOK-2 and a related human FasL clone, NOK-1 [Cat. No. 556371 (NA/LE)] may give different profiles in neutralization assays. It is thought that NOK-1 and NOK-2 likely recognize different FasL epitopes.

Flow Cytometry: Another clone, NOK-1, (556372, 556373) are recommended for flow cytometry. NOK-2 appears weaker for flow cytometry than NOK-1.

Western Blot: Neither NOK-1 nor NOK-2 are suggested for western blot analysis. Another human FasL clone, G247-4 (Cat. No. 556387) is suggested for the western blot analysis.

Suggested Companion Products

Catalog Number	Name	Size	Clone
553454	Purified Mouse IgG2a κ Isotype Control	0.5 mg	G155-178
556387	Purified Mouse Anti-Human CD178	0.1 mg	G247-4

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. An isotype control should be used at the same concentration as the antibody of interest.
3. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.

References

Kayagaki N, Kawasaki A, Ebata T, et al. Metalloproteinase-mediated release of human Fas ligand. *J Exp Med.* 1995; 182(6):1777-1783. (Clone-specific: ELISA, Immunoprecipitation, Neutralization)

Takahashi T, Tanaka M, Brannan CI, Jenkins NA, Copeland NG, Suda T, and Nagata S. Generalized lymphoproliferative disease in mice, caused by a point mutation in the Fas ligand. *Cell.* 1994; 76(6):969-976. (Biology)

Tanaka M, Suda T, Takahashi T, Nagata S. Expression of the functional soluble form of human Fas ligand in activated lymphocytes. *EMBO J.* 1995; 14(6):1129-1135. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

