## **Product Data Sheet**

## FITC anti-mouse I-Ad

Catalog # / Size: 1175030 / 500 µg

1175025 / 50 µg

Clone: 39-10-8

Isotype: Mouse IgG3, κ

(C3H x BALB/c)F<sub>1</sub> mouse cells Immunogen:

Reactivity: Mouse


**Preparation:** The antibody was purified by affinity

chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

**Concentration:** 0.5



BALB/c mouse splenocytes stained with 39-10-8 FITC

## **Applications:**

**Applications:** Flow Cytometry

Recommended

**Usage:** 


Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application** 

Notes:

Additional reported applications (for the relevant formats) include:

immunofluorescence microscopy2, and immunohistochemical staining of acetone-fixed frozen sections.



C57BL/6 mouse splenocytes stained with 39-10-8 FITC

**Application References:** 

- 1. Hiramine C, et al. 1995. Cell. Immunol. 160:157.
- 2. Wang Z, et al. 2004. J. Immunol. 172:5924.
  - 3. Ma XT, et al. 2006. Cancer Research 66:1169.
  - 4. Norian LA and Allen PM. 2004. J. Immunol. 173:835. PubMed
  - 5. Tian C, et al. 2007. J. Immunol. 179:6762.
  - 6. Ohashi K, et al. 2014. Mol Biol Cell. 25:828. PubMed

**Description:** 

The 39-10-8 antibody reacts with the I-Ad MHC class II alloantigen. These class II molecules are expressed on antigen presenting cells (including B cells) and a subset of T cells from H-2d bearing mice and are involved in antigen presentation to T cells expressing CD3/TCR and CD4 proteins. The 39-10-8 antibody does not cross-react with other haplotypes (a, b, k, p, q, s), but has been demonstrated to cross-react with the g7 haplotype.

**Antigen References:** 

- 1. Watts C. 1997. Ann. Rev. Immunol. 15:821.
- 2. Pamer E, et al. 1998. Ann. Rev. Immunol. 16:323.
  - 3. Wall KA, et al. 1983. J. Immunol. 131:1056.
  - 4. Ridgway WM, et al. 1998. J. E

